

Paper 8 Information Engineering Part A: Image Features and Matching  
**Solutions to Examples Paper**

1. *Images*

Each frame requires  $512 \times 512 \times 1 = 2.62 \times 10^5$  Bytes. A 25Hz stereo image stream requires  $2.62 \times 10^5 \times 25 \times 2 = 1.3 \times 10^7$  Bytes/s. Assuming an average A4 page of text contains 50 lines, with about 80 characters on each line, and that a character is represented (using an ASCII code) as a single byte, a page of text requires  $80 \times 50 \times 1 = 4000$  Bytes. So, instead of one second of stereo video, we could alternatively store  $1.3 \times 10^7 / 4000 \approx 3000$  pages of text — enough for a small encyclopaedia!

2. *Smoothing by convolution with a Gaussian*

Consider smoothing an image, first with a Gaussian of standard deviation  $\sigma_1$ , then with a Gaussian of standard deviation  $\sigma_2$ :

$$s(x) = g_{\sigma_2}(x) * (g_{\sigma_1}(x) * I(x))$$

Since convolution is associative, we can write this as the convolution of the image with the kernel  $g_{\sigma_2}(x) * g_{\sigma_1}(x)$ :

$$s(x) = (g_{\sigma_2}(x) * g_{\sigma_1}(x)) * I(x)$$

The easiest way to evaluate the convolution of two Gaussians is to find their Fourier transforms and then multiply the transforms in the frequency domain. If  $g_{\sigma}(x) \leftrightarrow G_{\sigma}(\omega)$ , then:

$$\begin{aligned} G_{\sigma}(\omega) &= \frac{1}{\sigma\sqrt{2\pi}} \int_{-\infty}^{\infty} \exp\left(-\frac{x^2}{2\sigma^2}\right) e^{-j\omega x} dx \\ &= \frac{1}{\sigma\sqrt{2\pi}} \int_{-\infty}^{\infty} \exp\left[-\left(\frac{x^2}{2\sigma^2} + j\omega x\right)\right] dx \\ &= \frac{1}{\sigma\sqrt{2\pi}} \int_{-\infty}^{\infty} \exp\left[-\frac{1}{2\sigma^2} (x^2 + 2j\omega\sigma^2 x)\right] dx \\ &= \frac{1}{\sigma\sqrt{2\pi}} \int_{-\infty}^{\infty} \exp\left[-\frac{1}{2\sigma^2} ((x + j\omega\sigma^2)^2 - j^2\omega^2\sigma^4)\right] dx \\ &= \exp\left(-\frac{\omega^2\sigma^2}{2}\right) \times \frac{1}{\sigma\sqrt{2\pi}} \int_{-\infty}^{\infty} \exp\left(-\frac{(x + j\omega\sigma^2)^2}{2\sigma^2}\right) dx \\ &= \exp\left(-\frac{\omega^2\sigma^2}{2}\right) \quad (\text{since the integral is a standard Gaussian}) \end{aligned}$$

Hence

$$\begin{aligned}
g_{\sigma_2}(x) * g_{\sigma_1}(x) &\leftrightarrow G_{\sigma_2}(\omega) \times G_{\sigma_1}(\omega) = \exp\left(-\frac{\omega^2 \sigma_2^2}{2}\right) \times \exp\left(-\frac{\omega^2 \sigma_1^2}{2}\right) \\
&\Leftrightarrow g_{\sigma_2}(x) * g_{\sigma_1}(x) \leftrightarrow \exp\left(-\frac{\omega^2(\sigma_2^2 + \sigma_1^2)}{2}\right)
\end{aligned}$$

The expression on the right is the Fourier transforms of a Gaussian with standard deviation  $\sqrt{\sigma_2^2 + \sigma_1^2}$ . So the convolution of two Gaussians with variances  $\sigma_1^2$  and  $\sigma_2^2$  is a Gaussian with variance  $\sigma_1^2 + \sigma_2^2$ . It follows that consecutive smoothing with a series of 1D Gaussians, each with a particular standard deviation  $\sigma_i$ , is equivalent to a single convolution with a Gaussian of variance  $\sum_i \sigma_i^2$ .

### Spatial domain convolution

Alternatively, we can convolve in the spatial domain. The trick, once again, is to complete the square:

$$\begin{aligned}
g_{\sigma_2}(x) * g_{\sigma_1}(x) &= \frac{1}{2\pi\sigma_1\sigma_2} \int_{-\infty}^{\infty} \exp\left(-\frac{u^2}{2\sigma_2^2}\right) \exp\left(-\frac{(x-u)^2}{2\sigma_1^2}\right) du \\
&= \frac{1}{2\pi\sigma_1\sigma_2} \int_{-\infty}^{\infty} \exp\left(\frac{-u^2\sigma_1^2 - x^2\sigma_2^2 - u^2\sigma_2^2 + 2ux\sigma_2^2}{2\sigma_1^2\sigma_2^2}\right) du \\
&= \frac{1}{2\pi\sigma_1\sigma_2} \int_{-\infty}^{\infty} \exp\left(\frac{-(\sigma_1^2 + \sigma_2^2)\left(u - \frac{x\sigma_2^2}{\sigma_1^2 + \sigma_2^2}\right)^2 + \frac{x^2\sigma_2^4}{\sigma_1^2 + \sigma_2^2} - x^2\sigma_2^2}{2\sigma_1^2\sigma_2^2}\right) du \\
&= \frac{1}{2\pi\sigma_1\sigma_2} \int_{-\infty}^{\infty} \exp\left(\frac{-\left(u - \frac{x\sigma_2^2}{\sigma_1^2 + \sigma_2^2}\right)^2}{\frac{2\sigma_1^2\sigma_2^2}{\sigma_1^2 + \sigma_2^2}}\right) \exp\left(\frac{-x^2\sigma_1^2\sigma_2^2}{2(\sigma_1^2 + \sigma_2^2)\sigma_1^2\sigma_2^2}\right) du \\
&= \frac{1}{2\pi\sigma_1\sigma_2} \exp\left(\frac{-x^2}{2(\sigma_1^2 + \sigma_2^2)}\right) \int_{-\infty}^{\infty} \exp\left(\frac{-\left(u - \frac{x\sigma_2^2}{\sigma_1^2 + \sigma_2^2}\right)^2}{2\left(\frac{\sigma_1\sigma_2}{\sqrt{\sigma_1^2 + \sigma_2^2}}\right)^2}\right) du \\
&= \frac{1}{\sqrt{2\pi}\sqrt{\sigma_1^2 + \sigma_2^2}} \exp\left(\frac{-x^2}{2(\sigma_1^2 + \sigma_2^2)}\right) \frac{1}{\sqrt{2\pi}\left(\frac{\sigma_1\sigma_2}{\sqrt{\sigma_1^2 + \sigma_2^2}}\right)} \int_{-\infty}^{\infty} \exp\left(\frac{-\left(u - \frac{x\sigma_2^2}{\sigma_1^2 + \sigma_2^2}\right)^2}{2\left(\frac{\sigma_1\sigma_2}{\sqrt{\sigma_1^2 + \sigma_2^2}}\right)^2}\right) du \\
&= \frac{1}{\sqrt{2\pi}\sqrt{\sigma_1^2 + \sigma_2^2}} \exp\left(\frac{-x^2}{2(\sigma_1^2 + \sigma_2^2)}\right) \quad (\text{since the integral is a standard Gaussian})
\end{aligned}$$

This expression is a Gaussian with standard deviation  $\sqrt{\sigma_2^2 + \sigma_1^2}$ .

### 3. Generating the Gaussian filter kernel

In general, if we discard the sample  $(n + 1)$  pixels from the center of the kernel, the size of the kernel will be  $2n + 1$  pixels. We can find  $n$  by solving:

$$\exp\left[-\frac{(n+1)^2}{2\sigma^2}\right] < \frac{1}{1000}$$
$$\Leftrightarrow n > 3.7\sigma - 1$$

So  $n$  must be the nearest integer to  $3.7\sigma - 0.5$ .

(a) Applying this formula for  $\sigma = 1$  gives  $n = 3$  and a kernel size of  $2n + 1 = 7$  pixels. The filter coefficients can be found by sampling the 1D Gaussian  $g_1(x)$  at the points  $x = \{-3, -2, -1, 0, 1, 2, 3\}$ . The sum of the coefficients is one, so regions of uniform intensity are unaffected by smoothing.

(b) For  $\sigma = 5$  we get  $n = 18$  and a kernel size of 37 pixels.

(c) The choice of  $\sigma$  depends on the *scale* at which the image is to be analysed. Modest smoothing (a Gaussian kernel with small  $\sigma$ ) brings out edges at a fine scale. More smoothing (larger  $\sigma$ ) identifies edges at larger scales, suppressing the finer detail. There is no right or wrong size for the kernel: it all depends on the scale we're interested in. Another factor is image noise: the smoothing suppresses noise. It may be difficult to detect fine scale edges, since a kernel large enough to suppress the noise may also suppress the fine detail. Finally, computation time may be an issue: large  $\sigma$  means a large kernel and computationally expensive convolutions.

### 4. Discrete convolution

The image and filter kernels are discrete quantities and convolutions are performed as truncated summations:

$$s(x) = \sum_{u=-n}^n g_\sigma(u)I(x-u)$$

Applying this to the pixel with intensity 118, which is the 11th pixel in the row, we obtain

$$\begin{aligned} s(x) &= \sum_{u=-3}^3 g_\sigma(u)I(11-u) \\ &= 0.004 \times 57 + 0.054 \times 77 + 0.242 \times 99 + 0.399 \times 118 \dots \\ &\quad + 0.242 \times 130 + 0.054 \times 133 + 0.004 \times 134 \\ &= 115 \quad (\text{to the nearest integer}) \end{aligned}$$

### 5. Differentiation and 1D edge detection

An approximation to the first-order spatial derivative of  $I(x)$  mid-way between the  $(n - 1)$  and  $(n + 1)$  sample is  $0.5(I(n + 1) - I(n - 1))$ . This can be computed by

convolving with the kernel 

1/2	0	-1/2
-----	---	------

 (remember that the kernel is flipped before the multiply and accumulate operation).

Applying this kernel to the smoothed row of pixels gives the approximation to the first-order spatial derivative:

x	x	x	x	2.5	3	5.5	11.5	17	18	14	8.5	3.5	0.5	-0.5	x	x	x	x
---	---	---	---	-----	---	-----	------	----	----	----	-----	-----	-----	------	---	---	---	---

The intensity discontinuity is at the maximum of the first-order spatial derivative. The maximum derivative (18) occurs at the tenth pixel - between the pixel with smoothed intensity 79 and the pixel with intensity 98<sup>1</sup>.

### 6. *Decomposition of 2D convolution*

The 2D convolution can be decomposed into two 1D convolutions as follows:

$$\begin{aligned}
 G_\sigma(x, y) * I(x, y) &= \frac{1}{2\pi\sigma^2} \int \int I(x-u, y-v) \exp\left(-\frac{u^2+v^2}{2\sigma^2}\right) du dv \\
 &= \frac{1}{\sqrt{2\pi}\sigma} \int \exp\left(-\frac{u^2}{2\sigma^2}\right) \left[ \frac{1}{\sqrt{2\pi}\sigma} \int I(x-u, y-v) \exp\left(-\frac{v^2}{2\sigma^2}\right) dv \right] du \\
 &= \frac{1}{\sqrt{2\pi}\sigma} \int \exp\left(-\frac{u^2}{2\sigma^2}\right) [g_\sigma(y) * I(x-u, y)] du \\
 &= g_\sigma(x) * [g_\sigma(y) * I(x, y)]
 \end{aligned}$$

Performing two 1D convolutions is much quicker than performing a single 2D convolution. A discrete 1D convolution with a kernel of size  $n$  requires  $n$  multiply and add operations. A discrete 2D convolution with a kernel of size  $n \times n$  requires  $n^2$  multiply and add operations. The speed-up offered by decomposing the 2D convolution is  $n^2/2n = n/2$ .

### 7. *Correlation and Convolution*

Convolution involves a reflection. They are identical if the kernel is symmetric.

8. *Feature detection and scale space* - see handout 2 and cribs for Tripos IB Paper 8 (F) 2010-2021.
9. *Interest point and Keypoint descriptors* - See handout 2 and cribs for Tripos IB Paper 8 (F) 2010-2021 on SIFT and normalised cross-correlation.
10. *Matching keypoints* - See handout 2 and cribs for Tripos IB Paper 8 (F) 2010-2021

Roberto Cipolla  
April 2022

---

<sup>1</sup>If you want to be more precise, you can localise the discontinuity to sub-pixel accuracy by calculating the second order derivatives and then interpolating to find the zero-crossing.